

30 January 1995

Prepared by

J. F. FENNELL and J. L. ROEDER  
Space and Environment Technology Center  
Technology Operations  
The Aerospace Corporation

M. GRANDE  
Rutherford Appleton Laboratories  
Great Britain

B. WILKEN  
Max-Planck-Institute for Aeronomy  
Germany

Prepared for

SPACE AND MISSILE SYSTEMS CENTER  
AIR FORCE MATERIEL COMMAND  
2430 E. El Segundo Boulevard  
Los Angeles Air Force Base, CA 90245

NOTES: ALL THIS REPRODUCTION  
WILL BE IN BLACK AND  
WHITE


Engineering and Technology Group

GROUP QUALITY IMPROVED 5

This report was submitted by The Aerospace Corporation, El Segundo, CA 90245-4691, under Contract No. F04701-93-C-0094 with the Space and Missile Systems Center, 2430 E. El Segundo Blvd., Los Angeles Air Force Base, CA 90245. It was reviewed and approved for The Aerospace Corporation by A. B. Christensen, Principal Director, Space and Environment Technology Center.

This report has been reviewed by the Public Affairs Office (PAS) and is releasable to the National Technical Information Service (NTIS). At NTIS, it will be available to the general public, including foreign nationals.

This technical report has been reviewed and is approved for publication. Publication of this report does not constitute Air Force approval of the report's findings or conclusions. It is published only for the exchange and stimulation of ideas.

  
LESLIE BELSMA, Maj, USAF  
Project Officer

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 30 January 1995		3. REPORT TYPE AND DATES COVERED
4. TITLE AND SUBTITLE Oxygen Charge State Abundance in the Inner Magnetosphere			5. FUNDING NUMBERS  F04701-93-C-0094	
6. AUTHOR(S) Fennell, J.F.; Roeder, J. L.; Grande, M.; and Wilken, B.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The Aerospace Corporation Technology Operations El Segundo, CA 90245-4691			8. PERFORMING ORGANIZATION REPORT NUMBER  TR-95(5940)-4	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Space and Missile Systems Center Air Force Materiel Command 2430 E. El Segundo Blvd. Los Angeles Air Force Base, CA 90245			10. SPONSORING/MONITORING AGENCY REPORT NUMBER  SMC-TR-95-11	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT  Approved for public release; distribution unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words)  The CRRES Magnetospheric Ion Composition Sensor (MICS) was used to obtain the relative oxygen charge state abundances in the earth's inner magnetosphere ( $3 < L < 7.5$ ). These abundances were obtained for average and quiet ( $K_p \leq 2+$ , $ DST  < 11$ ) conditions. They are presented as normalized spectra and compared to the predictions of Spjeldvik and Fritz (1978). These comparisons show that the observed spectra agree best with the Spjeldvik and Fritz predictions that assume an $O^+$ source, at the high L boundary in their model, for oxygen with $Q \leq +3$ but agree better with their predictions for a solar wind source, $O^{+6}$ , for oxygen with $Q > +3$ on average. As expected, $O^+$ is the dominant oxygen ion at all times in the inner magnetosphere for energies $> 60$ keV/Q. During average conditions the $Q \geq +4$ ions have fluxes comparable to $O^{+3}$ in the $L = 4.5 - 7.5$ region and are about 50% of the $O^{+3}$ fluxes in the $L = 3 - 5$ regions. Thus, the average $O^{+3}$ fluxes appear to represent a transition state between ionospheric and solar wind source oxygen.				
14. SUBJECT TERMS Magnetosphere, Ring Current, Trapped Particles, Oxygen			15. NUMBER OF PAGES 11	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT	

## **Acknowledgments**

The MICS sensor was provided by a team from the Max-Planck-Institute for Aeronomy, with support from the Max-Planck-Gesellschaft zur Förderung der Wissenschaften and the Rutherford Appleton Laboratory. The digital processing unit was provided by The Aerospace Corporation.

## Introduction

Over the last two decades there have been numerous observations of magnetospheric ion composition, first at the lower energies ( $< 30$  keV/q) by the S3-3, SCATHA, DE-1 and other satellites and later at extended energies by the AMPTE CCE (see Gloeckler et al., 1987 and references therein). These results focused on the major ion species  $H^+$ ,  $He^+$ ,  $He^{++}$  and  $O^+$ . More recently there have been reports on the higher charge states of oxygen and  $M > 4$  ions (Christon et al., 1994) in the outer magnetosphere. These have complemented the extensive results obtained in the interplanetary medium (Gloeckler and Geiss, 1989), in the magnetosheath (Gloeckler et al., 1986) and preliminary results in the magnetosphere (Kremser et al., 1985 and 1987).

The Christon et al. (1994) results provide a reference data set and boundary population for the transport of all ions into the inner magnetosphere. While several models and calculations have focused on describing the transport of the major ion species inward to form the ring current (Cornwall, 1972; Spjeldvik and Fritz, 1978; Sheldon and Hamilton, 1993; Chen et al, 1993, Sheldon, 1994) only Spjeldvik and Fritz, 1978 (hereafter denoted as S & F) have attempted to describe the distribution of oxygen ions. At the time S & F were developing their model only ion mass-composition was available for energetic oxygen ( $E_O > 30$  keV). Yet, in an attempt to match the observed oxygen ion spectra, S & F generated a model which predicted the oxygen charge state distributions for a wide range of energies in the inner magnetosphere and ring current for the two major oxygen sources, solar wind ( $O^{+6}$ ) and ionosphere ( $O^+$ ), as separate calculations. In this report we will provide the first observations of oxygen charge state spectra in the inner magnetosphere and will compare the observations with the predictions of S & F.

## Observations

For this study we used the ion charge state determination capabilities of the Magnetospheric Ion Composition Sensor (MICS) on the CRRES satellite (Wilken et al., 1992) to obtain the oxygen charge state spectra. The MICS sensor measures ions using time-of-flight (TOF), energy-per-charge ( $E/q$ ) and total energy ( $E$ ) measurements for each ion detected (Wilken et al., 1992). The energy per charge measured ranges from 1 to 425 keV/q in 32 logarithmic steps. The parameters  $E$ , TOF and  $E/q$  are used to determine the energy, mass and charge state of the ions. To obtain the relative oxygen intensity at each charge state we used the highest resolution three-parameter or DE data where the TOF,  $E/q$  and  $E$  are transmitted for a subset of ions measured ( $\sim 4$  events/sec). Because of the limited sampling rates one cannot use the DE data to obtain absolute fluxes. What the DE data do provide, in all instrument modes, is the relative fluxes of ions with different charge states but the same mass. All the MICS DE data from the period 12 January through 17 September 1991 were extracted and tagged with appropriate ephemeris parameters. These data were then ordered by the values of their three parameters and accumulated into arrays of size  $256 \times 1024 \times 32$  for each half of a CRRES orbit into 21 bins in

L (Roeder et al., 1994). The resultant data base was then sampled in L and according to an activity criterion. First all data from orbits 416-1001 were accumulated into L bins covering the L ranges 3 - 5 and 4.5 - 7.5 for all times to obtain all oxygen ion intensities for average conditions. (Orbits prior to 416 were not used because of the operational mode of the instrument.) Separately, the quiet-time oxygen ion data only were accumulated into the same L bins. The quiet time was defined according to a criterion used by Sheldon and Hamilton (1992) and Roeder et al. (1994). To be considered quiet, the  $|DST|$  and  $K_p$  for a data sample had to be  $\leq 11$  and  $\leq 2+$  respectively and also  $\leq 16$  and  $\leq 3$  respectively for the previous 24 hours.

These DE data were then examined and the oxygen ions were extracted based on their known  $E$  versus TOF dependence (from calibration data). The oxygen ions were binned into ion energy per charge ( $E/q$ ) and ion total energy ( $E$ ) intervals ( $32 \times 256$  array). A multiple Gaussian fit, as shown in Figure 1, was applied to the resulting intensity versus  $E$  for each  $E/q$  interval separately. These fits allowed us to extract the relative intensities at the peak of each observed oxygen charge state as a function  $E/q$  as indicated by the labels in the example shown in Figure 1.




Figure 1 Average 175 keV/q oxygen intensity versus energy channel number for  $L > 4.5$  (square points). The solid curve is a fit to the data using a superposition of Gaussians. The labels on the peaks indicate the oxygen charge states.

These intensities were used to form a spectrum for each charge state. The resultant spectra were normalized to the intensity of the 100 keV/Q  $O^+$  ions. The intensities of the higher charge states ( $+2 \leq Q \leq +7$ ) represent the relative abundances of these ions compared to  $O^+$ . Figure 2 shows the resultant normalized oxygen intensities versus energy for all magnetic conditions (top panels) and for quiet conditions (bottom panels) for  $4.5 < L < 7$  (right panels) and  $3 < L < 5$  (left panels). These we call the oxygen charge state spectra. These data were averaged over all local times except the interval from 4 to 10 MLT on the morning and pre-noon side and cover the period from 12 January through 17 September 1991.

As was expected and can easily be seen in Figure 2,  $O^+$  is the dominant oxygen ion at all energies in both regions of L for the average and magnetically quiet conditions. The average  $K_p$  and  $D_{ST}$  for this time period were 3 and -26.7, respectively. The total duration of the quiet intervals was too short to obtain statistically significant counting rates for the higher charge states of oxygen. In the quiet time  $3 < L < 5$  region no measure of  $O^{\geq +4}$  intensities were obtained, except for a single  $O^{+4}$  intensity value at 1500 keV (triangle point in the lower left panel of Figure 2). At higher L the quiet time  $O^{+4}$  was well measured, but the  $O^{+5}$  was poorly measured and only a single point for  $O^{+6}$  was obtained (diamond point near 300 keV in the lower right panel of Figure 2).

We then extracted the relative oxygen ion fluxes at 100 - 200 keV and compared their levels with the predictions of S & F as shown in Figure 3. The MICS  $O^+$  fluxes have been normalized to the S & F fluxes. The S & F curves were generated assuming that  $O^+$  was the source species for oxygen (from S & F's Figure 8). At the lower L the agreement with the first low charge states of oxygen ( $O^{++}$  and  $O^{+++}$ ) is quite good both for the average and quiet conditions. The higher L and higher charge state values are not well represented by the F & S calculation. Still, this is quite good agreement considering the limited source assumption and the fact that S & F did not have charge state information as a reference for their calculations.

Comparison of the MICS oxygen spectra with the predictions of S & F (from their Figure 8) is shown in Figure 4. Again, the MICS  $O^+$  data is normalized to the S & F predictions for  $O^+$  at 100 keV. One sees, like in Figure 3, that the low oxygen charge state spectra agree best with these S & F curves for both the average and quiet conditions. At low L the MICS data and the S & F model predictions agree best for energies below ~200 keV for both average and quiet conditions. At the higher energies the observed spectra are harder than predicted. Also, the higher charge state spectra are clustered closer together than predicted. This is expected because we are comparing to the S & F prediction based on an assumed  $O^+$  source for the oxygen. Examination of the S & F predicted spectra for an assumed solar wind source ( $O^{+6}$ , their Figure 9) shows that they would have the spectra for all the oxygen charge states clustered closer together.

Similarly, at high L the shapes of the observed  $O^+$  spectra for average and quiet conditions are close to the S & F  $O^+$  spectrum, but the observed spectral shapes for higher charge states are somewhat steeper and closer together. The S & F predictions for the solar wind source would have all the different charge state spectra clustered closer together but with slopes similar to their  $O^+$  source curves. Still, the agreement is reasonably good given the lack of charge state information at the time of their work.

## Discussion

A careful examination of Figure 2 shows that, in general, the spectra for  $O^{++}$  is consistent with being derived from  $O^+$  via charge exchange. This is expected because the charge exchange cross sections  $\sigma_{12} \sim \sigma_{21}$ , in the notation of S & F, for the range of energies covered there (see Figures 3 and 4 from S & F). For  $O^{+3}$  the picture is more complicated, as shown in S & F. The cross section for electron capture (charge loss)  $\sigma_{32}$  is a decreasing function of energy, whereas that for electron loss,  $\sigma_{23}$ , (charge gain) is an increasing function of energy with  $\sigma_{32} \sim \sigma_{23}$  near 10<sup>3</sup> keV.

At the lower L range,  $O^{+3}$  is most likely the transition charge state with comparable contributions from charge loss and charge gain processes from adjacent charge states. For  $L \geq 4.5$ , under average conditions (top right panel of Figure 2), the  $O^{+4}$  intensities are comparable to  $O^{+3}$  and those for  $O^{+5}$  and  $O^{+6}$  are higher. Thus, one may argue that most of the  $O^{+3}$  at the higher L must derive from oxygen with charge states  $\geq +4$  via charge exchange because  $\sigma_{34} < \sigma_{43}$  below ~ 1500 keV. However, it is clear from the relative intensities of the  $O^{+3}$  in the other panels that  $O^{+3}$  is also derived predominantly from  $O^{+2}$  via upward charge exchange. Thus  $O^{+3}$  below 1000 keV in energy is a transition charge state separating the oxygen ions that are dominated by the ionosphere as a source from those dominated by the solar wind as a source. What is unclear is which source is dominant for the higher energy oxygen ions.

S & F did well given the limited input data that they had to work with. We know better now, based on the results of Christon et al. (1994), what the real source spectrum for oxygen is and what combination of ionospheric and solar wind ions is required. It is time to utilize these data to redo the S & F calculations. It is clear that such a calculation should be able to match the CRRES MICS data taken in the inner magnetosphere very well.




Figure 2 Oxygen charge state spectra for average (top panels) and quiet (bottom panels) conditions and for  $L = 3 - 5$  (left panels) and  $L = 4.5 - 7.5$  (right panels). The energy spectra of the different charge states are color coded as noted in the legends. The spectra are normalized to the  $O^+$  data at 100 keV in each panel.




Figure 3 Comparison of CRRES MICS 100 - 200 keV ion intensities for different oxygen charge states with the predictions of Spjeldvik and Fritz (1987) for 100 keV oxygen. CRRES data from two different L ranges and for average (left panel) and quiet (right panel) magnetic conditions. The Spjeldvik and Fritz curves were based on the assumption that  $O^+$  was the only ion in the source region at high L shells ( $L > 7$ ). The CRRES results for  $O^+$  are normalized to the Spjeldvik and Fritz results for  $O^+$ .






Figure 4 Comparison of CRRES MICS oxygen charge state spectra with the predictions of Spjeldvik and Fritz (1978) for an assumed high L boundary source composed of only O<sup>+</sup>. The data are for average magnetic conditions (top panels), quiet times (bottom panels). The MICS data from L = 3 - 5 is compared to S & F results for L=4 (left panels) and MICS data from L = 4.5 - 7.5 is compared to the S & F results for L = 6.1 (right panels).

## References

- Chen, M., M. Schulz, L. R. Lyons, and D. J. Gorney, Stormtime transport of ring current and radiation-belt ions, *J. Geophys. Res.*, 99, 3835-3849, 1993.
- Christon, S. P., D. C. Hamilton, G. Gloeckler, T. E. Eastman, F. M. Ipavich, High charge state carbon and oxygen ions in Earth's equatorial quasi-trapping region, *J. Geophys. Res.*, 99, 13465-13488, 1994.
- Cornwall, J. M., Radial diffusion of ionized helium and protons: A probe for magnetospheric dynamics, *J. Geophys. Res.*, 77, 1756-1770, 1972.
- Gloeckler, G. and D. C. Hamilton, AMPTE Ion Composition Results, *Physica Scripta*, T18, 73-84, 1987.
- Gloeckler, G., and J. Geiss, The abundances of elements and isotopes in the solar wind, in *Proceedings of the Symposium on Cosmic Abundances of Matter*, edited by C. Waddington, Conf. Proc. 183, pp. 49-71, American Institute of Physics, New York, 1989.
- Gloeckler, G., F. M. Ipavich, D. C. Hamilton, B. Wilken, W. Studemann, G. Kremser, and D. Hovestadt, Solar wind carbon, nitrogen and oxygen abundances measured in the Earth's magnetosheath with AMPTE/CCE, *Geophys. Res. Lett.*, 13, 793-796, 1986.
- Kistler, L. M., F. M. Ipavich, D. C. Hamilton, G. Gloeckler, B. Wilken, G. Kremser, and W. Studemann, Energy spectra of the major ion species in the ring current during geomagnetic storms, *J. Geophys. Res.*, 94, 3579-3599, 1989.
- Kremser, G., W. Studemann, B. Wilken, G. Gloeckler, D. C. Hamilton and F. M. Ipavich, Average spatial distributions of energetic  $O^+$ ,  $O^{2+}$ ,  $O^{6+}$ , and  $C^{6+}$  ions in the magnetosphere observed by AMPTE CCE, *J. Geophys. Res.*, 92, 4459-4466, 1987.
- Kremser, G., W. Studemann, B. Wilken, G. Gloeckler, D. C. Hamilton, F. M. Ipavich and D. Hovestadt, Charge state distributions of oxygen and carbon in the energy range 1 to 300 keV/e observed with AMPTE/CCE in the magnetosphere, *Geophys. Res. Letters*, 12, 847-850, 1985.
- Roeder, J. L., J F Fennell, M W Chen, M Schulz, M Grande and S Livi, CRRES observations of the composition of the ring current ion populations, *Adv. in Sp. Res.*, in press, 1995.
- Sheldon, R. B., and D. C. Hamilton, Ion transport and loss in the Earth's quiet time ring current, 1. Data and standard model, *J. Geophys. Res.*, 98, 13491-13508, 1993.
- Sheldon, R.B., Ion transport and loss in the Earth's quiet time ring current, 2. Diffusion and magnetosphere-ionosphere coupling, *J. Geophys. Res.*, 99, 5705-5720, 1994.
- Spjeldvik, W. N., and T. A. Fritz, Theory for charge states of energetic oxygen ions in the Earth's radiation belts, *Journal of Geophys. Res.*, 83, 1583-1594, 1978.
- Spjeldvik, W. N., Expected charge states of energetic ions in the magnetosphere, *Space Sci. Rev.*, 23, 499-538, 1979.
- Wilken, B., W. Weiss, D. Hall, M. Grande, F. Soraas, J. F. Fennell, Magnetospheric ion composition spectrometer onboard the CRRES spacecraft, *J. Spacecraft and Rockets*, 29, 585-591, 1992.